

Title

The Relationship Between Cannabis Legalization, Suicide and Mental Health Remains Unchanged After Recreational Access

Jeffrey A. Singer, MD¹, Jacob James Rich, MA², Michael Schemenaur, MA³, and Robert Capodilupo, MPhil⁴

1. Health Policy Studies, Cato Institute, Washington, DC
2. Drug Policy Project, Reason Foundation, Washington, DC
3. Department of Economics, Eller College of Management, University of Arizona, Tucson, AZ
4. Yale Law School, New Haven, CT

Abstract:

Background. We extend the model introduced by Anderson et al. [M.D. Anderson, D.I. Rees, J.J. Sabia, *American Journal of Public Health* 104, 2369-2376] to evaluate the public mental health implications of new developments in marijuana policy, such as recreational marijuana access and additional years of data on suicide mortality.

Methods. We obtained state-level suicide data from the National Vital Statistics System's Mortality Detail Files for 1999-2018. We used panel regression analysis to examine the association between suicides per 100 000 population and both medical and recreational marijuana access. Three specifications of the equation were employed as a robustness test.

Results. After adjusting for economic conditions, alcohol taxes and consumption, time effects, state fixed effects, and state-specific linear time trends, the legalization of recreational marijuana was associated with a 2.9% (95% confidence interval [CI] = -5.6, -.03) reduction in the suicide rate for the most rigorous specification, but didn't pass the robustness check at the $P < .05$ level for the other two specifications. However, recreational marijuana legalization was associated with a 5.4% reduction (95% CI = -9.4%, -1.4%) in suicide rates for males in the 40 to 49 age group, passing the robustness check for all specifications.

Conclusion. Suicide rates among men in the 40 to 49 age group fell after recreational marijuana legalization compared with those in states that did not legalize, which is consistent with the results found by Anderson et al. for medical marijuana when that same population was younger. There was no consistent relationship between medical marijuana legalization and suicide rates for any population. Additionally, mental health indicators at the state level were unaffected by either recreational or medical marijuana legalization. We confirm the findings of Anderson et al. (2014), observing no consistent relationship between marijuana legalization and observable mental health outcomes. However, suicide rates increase at lower rates on average relative to other states following recreational marijuana legalization.

1 **The Relationship Between Cannabis Legalization, Suicide and Mental Health Remains** 2 **Unchanged After Recreational Access**

3
4 As of September 1 2020, 11 states and the District of Columbia have voted to legalize marijuana
5 for recreational use, while medical marijuana is available in 33 states. A November 2019 survey
6 from the Pew Research Center found that 67 percent of adults support national legalization of
7 marijuana, including 63 percent of the Baby Boomer generation, 76 percent of Millennials, and
8 65 percent of Gen X. (Daniller, 2019) In October 2018 Canada became the first G7 country to
9 federally legalize marijuana for recreational use, and the same month Mexico's Supreme Court
10 ruled marijuana prohibition unconstitutional, setting the stage for legalization in that country.
11 (Hudak, 2020) In the not-too-distant future marijuana use in the United States might be federally
12 decriminalized and the matter devolved to the states. Legislation to that effect was introduced in
13 the 116th Congress. (Nadler, 2020) These developments reflect the increasing acceptance of the
14 recreational use of marijuana, which trends suggest will further increase in days ahead.

15 Policy makers considering marijuana legalization should consider the public health
16 implications of increasing access to a psychoactive substance that could lead to substance use
17 disorder. In a 2009 Norwegian study, Pedersen concluded “exposure to marijuana by itself does
18 not lead to depression but that it may be associated with later suicidal thoughts and attempts.”
19 (Pederson, 2008) A 2014 systematic review and meta-analysis of longitudinal studies by
20 Canadian researchers found an increased risk for developing depressive disorders may be
21 associated with heavy marijuana use. (Lev-Ran, Roerecke, George, McKenzie, & Rehm, 2014)
22 Agrawal et al., in a 2017 retrospective study, found that the correlation between marijuana use,
23 suicidal thoughts, and major depressive disorders in discordant twins was not solely attributable
24 to predisposing factors. (Agrawal, et al., 2017) This observation confirmed similar findings
25 reported by researchers at the same institution ten years earlier. (Lynskey, et al., 2004) A 2015
26 study using community-based samples from the Australian Twin Registry by Delforterie et al.
27 found a “modest association” of marijuana use with suicidal thoughts and attempts, concluding
28 that the association of suicidal thoughts and behaviour with marijuana use disorder requires
29 further study. (Delforterie, et al., 2015) In a 2020 study in *JAMA Network Open*, Gorfinkel et al.
30 found that people reporting depression tend to use marijuana at a higher rate. (Gorfinkel, Stohl,
31 & Hasin, 2020)

32 However, much of the literature finds no relationship between marijuana use and mental
33 health outcomes. In particular a 2014 regression analysis of state-level data from the National
34 Vital Statistics System (NVSS) Mortality Detail Files from 1990-2007 by Anderson et al. found
35 no general association between medical marijuana legalization and suicide rates, but found
36 10.8% and 9.4% reductions in the suicide rates in males age 20 to 29 and 30 to 39 in the U.S.,
37 respectively. (Anderson, Rees, & Sabia, 2014) Additionally, a 2018 Canadian study found no
38 correlation between suicidal behaviour and heavy marijuana use in male or female patients with
39 psychiatric disorders. (Naji, et al., 2018) With the mental outcomes of marijuana use contested
40 by much of the literature, policymakers would benefit from research investigating the population
41 effects on mental health after marijuana legalization.

42 Marijuana use is tracked by the Substance Abuse and Mental Health Services
43 Administration (SAMHSA) with the National Survey on Drug Use and Health (NSDUH). As of
44 2018, about 43.5 million people (or 15.9 percent of the U.S. population above the age of 12)
45 consumed marijuana at least once that year—a historic high. Of those marijuana users, 3.1
46 million used marijuana for the first time, which has been a relatively consistent finding every
47 year since Colorado became the first state to legalize marijuana in 2014. That year 12.9 percent
48 of the U.S. population used marijuana in the past year and the recent trend of increase is
49 predicted to continue into the near future. (SAMHSA, 2015) Marijuana use reported in the past
50 month has also steadily increased in the U.S. since 2007, after a subtler increase starting after an
51 all-time low in 1992. (Golub, Johnson, & Dunlap, 2005)

52 Reports of major depressive episodes have increased since 2010, while both serious
53 thoughts of suicide and severe mental illness have increased each year since the inception of the
54 survey questions in 2008. (SAMHSA, 2019) Suicide rates in the U.S. have been steadily
55 increasing since an all-time low in 1999 and were at an all-time high in 2018. Respondents who
56 reported past month marijuana use in 2017 were much more likely to experience major
57 depression (2.29 times), suicidal thoughts (3.2 times), and slightly more prone to any/severe
58 mental illness (1.16/1.64 times) than nonusers. (SAMHSA, 2018) This observation has garnered
59 much interest in the relationship between marijuana use and possible mental health deterioration.
60 But with increasing evidence of marijuana's medicinal benefits, and with its growing popularity
61 as a recreational drug, it is necessary to definitively answer the questions surrounding heavy
62 marijuana use and risk of depression and suicide.

63 In the years since the Anderson et al. review of data from 1990-2007, the advent of
64 recreational marijuana laws has provided unprecedented access to marijuana. Significant gaps
65 remain in the knowledge of the potential harms and benefits of marijuana use, and the effects
66 these laws have on various facets of mental illness are a fertile topic for new research. (National
67 Academies of Sciences, Engineering, and Medicine, 2017)

68

69 **Methods**

70

71 Employing most of the same techniques used by Anderson et al. (2014), we use population-
72 weighted ordinary least squares (OLS) regressions for the observed groups to conduct a state-
73 level panel analysis for various time periods of available data to estimate the effect recreational
74 and medical marijuana laws have had on various proxies for mental health: suicide rates per 100
75 000 people for the general population, males, females, and both sexes separately for the age
76 groups 15-19, 20-29, 30-39, 40-49, 50-59, and 60 and above (1999-2018); the percentages of the
77 population that reported “serious thoughts of suicide in the past year,” “any mental illness in the
78 past year,” and “serious mental illness in the past year” for the age groups 18-25, 18 and up, and
79 26 and up (2009-2018); and the percentages of the population that reported they “had at least one
80 major depressive episode in the past year” for the age groups 12-17, 18-25, 18 and up, and 26
81 and up (2006-2018).

82 Similar to Anderson et al. (2014), we control for observable state-level characteristics the
83 literature considers risk factors for suicide: unemployment rates, GDP per capita, beer taxes, and
84 ethanol consumption per capita for the population age 21 and older. Also following Anderson et
85 al. (2014), we attempt to control for unobserved factors with time effects, fixed effects, and state-
86 specific linear time trends. Since previous studies suggest stricter alcohol policies can reduce
87 suicides, (Markowitz, Chatterji, & Kaestner, 2003; Birckmayer & Hemenway, 1999; Jones,
88 Pieper, & Robertson, 1992; Robertson, 2014) Anderson et al. (2014) controlled for whether a
89 0.08 blood-alcohol-content (BAC) law was in effect, whether a zero-tolerance drunk-driving law
90 was in effect, and for the beer excise tax. We removed the controls for BAC laws because every
91 state had implemented a 0.08 BAC limit for our entire period, except for Utah, which
92 implemented a .05 limit December 25 2018. Additionally, we removed indicators for zero-
93 tolerance drunk-driving laws because every state had implemented such restrictions by 2001.

94 (Carpenter, Heavy alcohol use and youth suicide: Evidence from tougher drunk driving laws,
95 2004; Carpenter, How do Zero Tolerance Drunk Driving Laws work?, 2004) We argue those
96 variables ultimately attempted to capture state government interventions that reduce alcohol use,
97 which is why we added a measurement for per capita ethanol consumption.

98 To test the robustness of our findings, we follow Anderson et al. (2014) and employ a
99 sensitivity analysis involving three separate specifications for each dependent variable: first only
100 adjusting for state and year effects; then adjusting for state effects, year effects, and covariates;
101 and finally adjusting for state effects, year effects, covariates, and state-specific linear time
102 trends in the most rigorous model. Robust standard errors were employed and clustered at the
103 state level for all specifications. Additionally, we took the natural log transform of all dependent
104 variables, and because the log of zero is undefined, we dropped all observations of zero suicides.
105 If the coefficients for the marijuana law indicators remain relatively stable and significant for all
106 regression specifications, there is evidence that a predictive relationship exists between
107 marijuana legalization and the mental health variables of interest. (Miron & Tetelbaum, 2009;
108 Thornton, 2011)

109 Suicide mortality (X60-X84, Y87.0, and U03 intentional self-harm) data and population
110 estimates are sourced from the Centers for Disease Control and Prevention (CDC) Wide-ranging
111 Online Data for Epidemiologic Research (WONDER) NVSS database, reports of mental illness
112 from the SAMHSA NSDUH, unemployment rates and GDP per capita from the Bureau of Labor
113 Statistics, ethanol consumption from the National Institute on Alcohol Abuse and Alcoholism,
114 and beer excise tax figures from the Tax Policy Center at the Urban Institute and Brookings
115 Institution. Marijuana policy indicators were calculated referencing reports by the Marijuana
116 Policy Project and the National Organization for the Reform of Marijuana Laws (NORML).
117 These indicators consider the dates laws were enacted and for the first year assume a value
118 between 0 and 1 equal to the proportion of days the policies were effective. The year recreational
119 marijuana laws became effective were used to graph unadjusted suicides per 100 000 population
120 over the year of law change to compare rates with states that did not legalize recreational
121 marijuana anytime during the period.

122

123 **Results**

124

125 Table 1 presents the dates that various marijuana laws went into effect. Average suicide rates per
126 100 000 population over the year of law change are displayed in Figure 1, graphing average
127 suicides rates for the five jurisdictions that have legalized recreational marijuana for at least three
128 years (Alaska, Colorado, District of Columbia, Oregon, and Washington) next to the 40 states
129 that didn't legalize marijuana for the entire period of interest centred at 2014, the average year of
130 implementation for the 5 legalizing states. Reviewing the slopes of both groups before policies
131 went into effect, states that would eventually legalize recreational marijuana had a higher
132 average annual increase in suicide rates (0.708 per 100 000 people) relative to states that never
133 legalized (0.374 per 100 000 people). However, after legalization policies went into effect, states
134 permitting access to recreational marijuana reduced their average annual increase to a level
135 (0.339 per 100 000) below both groups before legalization and the non-legalizing group after
136 legalization (0.411 per 100 000).

137 Table 2 presents the regression results measuring the effect marijuana legalization had on
138 the mental health outcomes. The only relationship that passed each of the robustness check
139 specifications was between suicide rates for males in the 40 to 49 age group and recreational
140 marijuana legalization, where legalization is followed by a 5.4% reduction (95% confidence
141 interval [CI] = -9.4%, -1.4%) in suicides per capita in the most rigorous model. Interestingly,
142 Anderson et al. found that medical marijuana legalization led to 10.8% (95% CI = -17.1%,
143 -3.7%) and 9.4% (95% CI = -16.1%, -2.4%) drops in the 20 to 29 and 30 to 39 age groups,
144 respectively, many of whom would now be in the 40 to 49 age group. This population is now
145 older because the youngest data from the previous study were published 13 years ago.

146 The only other relationship that was significant for the most rigorous specification was a
147 2.9% reduction (95% CI = -5.6, -.03) in the total suicide rate following recreational marijuana
148 legalization. The relationship between recreational marijuana legalization was also weakly
149 associated reductions in the total suicide rate for the other two specifications, but the estimates
150 didn't quite achieve significance at the standard $P < .05$ level.

151

152 **Discussion**

153

154 Despite suicide rates reaching unprecedented levels across the country, suicides in states with
155 recreational marijuana access have increased at slower rates after their policies went into effect

156 relative to their rates before legalization and relative to other states across the country. Our
157 analysis finds strong evidence that recreational marijuana laws reduce suicide rates for middle
158 age men and weak evidence recreational marijuana laws reduce suicide rates for the population
159 as a whole. In line with the conclusions of Anderson et al. (2014), we find that marijuana laws in
160 general don't have consistent effects on suicide rates or reports of mental illness, despite
161 correlations between marijuana use and reports of depression.

162 While 24% of past year marijuana users report depression, 25.1% of people who report
163 depression admit to using marijuana in the past year. (SAMHSA, 2018) The correlations
164 concerning policy makers may be due to simultaneity, or "reverse causality," which could be the
165 result of self-medication. Regardless, there is no evidence that the liberalization of marijuana in
166 the U.S. is leading to reductions in public mental health.

167

168 **Limitations**

169

170 Unfortunately, due to the limitations of public SAMHSA data because of privacy
171 concerns, it is not possible to utilize geographical controls for individual data. Our analysis
172 suffers from the same limitations that plague observational studies in general. (Koopman &
173 Lynch, 1999) To improve on future public research independent of the government's efforts,
174 SAMHSA should add state-level identifiers to its public individual-response data for all years
175 after 2002. Since NSDUH state-level samples are too large to identify the individuals who took
176 the survey, privacy will not be a concern if state of residence information is released to the public
177 for each individual response. Permitting state-level identifiers will allow for the employment of
178 accepted metric methods that use geographic identifiers to control for environmental and
179 reporting differences. The consequences of drug use in general are of the utmost interests to
180 public health officials, and the literature would be best advanced by allowing public access to
181 currently restricted data.

182 There have also been legitimate challenges to the accuracy of the NSDUH and suicide
183 rates. Critics claim that because the NSDUH requires the surveyed individuals to have a
184 residence, much of the population (especially homeless people) are not represented by the
185 estimates. (Hedden, et al., 2012) Additionally, it is widely believed suicide rates are
186 underreported. (Warshauer & Monk, 1978) These are legitimate criticisms, but the regression

187 analyses are much more concerned with the changes in rates, as opposed to the absolute accuracy
188 of the rates themselves. Additionally, there's no reason to believe states with recreational
189 marijuana access would be more likely to underreport suicides. Since consistent methods for
190 collecting these data were employed for the entire period of interest for all dependent variables,
191 we are confident that the variable regression coefficients reflect the relationships between
192 evolving trends of substance availability and the mental health identifiers of interest.

193

194 **Public Health Implications and Conclusion**

195

196 The use of any drug, psychoactive or otherwise, entails certain risks along with any benefits.
197 Marijuana is no exception. Critics of marijuana legalization point to studies showing correlations
198 between heavy cannabis use and suicide, depression, and mental health disorders. However, such
199 studies demonstrating correlation have yet to confirm causation, which should be determined by
200 a model's ability to predict. (Arrow, 1963) Although those reporting depression to SAMHSA
201 have increasingly used marijuana since states began increasing access to regulated cannabis,
202 (Gorfinkel, Stohl, & Hasin, 2020) we observe no evidence that these laws have predictive
203 relationships with reports of any mental illness. Additionally, recreational marijuana access is
204 followed by reductions in suicide for some age groups. Anderson et al. (2014) had found similar
205 relationships between medical marijuana access and suicide rates, and the lack of an association
206 between mental health and medical marijuana access in our study is likely because recreational
207 marijuana access is a superior substitute requiring fewer barriers to purchase cannabis.

208 We propose as medicinal and recreational use of marijuana becomes more widespread
209 and mainstream, concerns about the correlation between marijuana use and depression shouldn't
210 interfere with state or federal efforts to decriminalize or legalize marijuana. In fact, legalization
211 will have the salutary effect of allowing more rigorous research—now inhibited by federal
212 prohibition—into the further benefits, as well as any other potential harms, from the long-term
213 use of marijuana, and promote safer use.

214 **References**

215

216 Agrawal, A., Nelson, E. C., Bucholz, K. K., Tillman, R., Grucza, R. A., Statham, D. J., . . .
217 Lynskey, M. T. (2017, September 1). Major depressive disorder, suicidal thoughts and

218 behaviours, and cannabis involvement in discordant twins: a retrospective cohort study.
219 *Lancet Psychiatry*, 4(9), 704-714.

220 Anderson, M. D., Rees, D. I., & Sabia, J. J. (2014, December). Medical Marijuana Laws and
221 Suicides by Gender and Age. *American Journal of Public Health*, 104(12), 2369-2376.

222 Arrow, K. J. (1963, December). Uncertainty and the Welfare Economics of Medical Care. *The*
223 *American Economic Review*, 53(5), 941-973.

224 Birckmayer, J., & Hemenway, D. (1999). Minimum-age drinking laws and youth suicide, 1970-
225 1990. *American Journal of Public Health*, 89(9), 1365-1368.

226 Bureau of Labor Statistics. (2020). Databases, Tables & Calculators by Subject. Washington,
227 District of Columbia, United States.

228 Carpenter, C. (2004, May). Heavy alcohol use and youth suicide: Evidence from tougher drunk
229 driving laws. *Journal of Policy Analysis and Management*, 23(4), 831-842.

230 Carpenter, C. (2004, January). How do Zero Tolerance Drunk Driving Laws work? *Journal of*
231 *Health Economics*, 23(1), 61-83.

232 Centers for Disease Control and Prevention. (2020). Wide-ranging Online Data for
233 Epidemiologic Research (WONDER). Atlanta, Georgia, United States.

234 Daniller, A. (2019, November 14). Two-thirds of Americans support marijuana legalization.
235 *News In The Numbers*.

236 Delforterie, M. J., Lynskey, M. T., Huijink, A. C., Creemers, H. E., Grant, J. D., Few, L. R., . . .
237 Agrawal, A. (2015, May 1). The relationship between cannabis involvement and suicidal
238 thoughts and behaviors. *Drug and Alcohol Dependence*, 150, 98-104.

239 Golub, A., Johnson, B. D., & Dunlap, E. (2005). The Growth in Marijuana Use Among
240 American Youths During the 1990s and the Extent of Blunt Smoking. *Journal of*
241 *Ethnicity in Substance Abuse*, 4(3-4), 121.

242 Gorfinkel, L. R., Stohl, M., & Hasin, D. (2020, August 18). Association of Depression With
243 Past-Month Cannabis Use Among US Adults Aged 20 to 59 Years, 2005 to 2016. *JAMA*
244 *Network Open*, 3(8).

245 Hedden, S., Gfroerer, J., Barker, P., Smith, S., Pemberton, M. R., Saavedra, L. M., . . . Novak, S.
246 P. (2012). *Comparison of NSDUH Mental Health Data and Methods with Other Data*
247 *Sources*. Rockville, MD: Department of Health and Human Services.

248 Hudak, J. (2020). *Marijuana: A Short History*. Washington, D.C.: Brookings Institution.

249 Jones, N. E., Pieper, C. F., & Robertson, L. S. (1992, January). The effect of legal drinking age
250 on fatal injuries of adolescents and young adults. *American Journal of Public Health*,
251 82(1), 112-115.

252 Koopman, J. S., & Lynch, J. W. (1999, October 7). Individual causal models and population
253 system models in epidemiology. *American Journal of Public Health*, 89(8), 1170-1174.

254 Lev-Ran, S., Roerecke, M., George, T. P., McKenzie, K., & Rehm, J. (2014, March). The
255 association between cannabis use and depression: a systematic review and meta-analysis
256 of longitudinal studies. *Psychological Medicine*, 44(4), 797-810.

257 Lynskey, M. R., Glowinski, A. L., Todorov, A. A., Bucholz, K. K., Madden, P. A., Nelson, E.
258 C., . . . Heath, A. C. (2004). Major Depressive Disorder, Suicidal Ideation, and Suicide
259 Attempt in Twins Discordant for Cannabis Dependence and Early-Onset Cannabis Use.
260 *JAMA Psychiatry*, 61(10), 1016-1032.

261 Markowitz, S., Chatterji, P., & Kaestner, R. (2003, March). Estimating the impact of alcohol
262 policies on youth suicides. *Journal of Mental Health Policy Economics*, 6(1), 37-46.

263 Miron, J. A., & Tetelbaum, E. (2009, April). Does the Minimum Legal Drinking Age Save
264 Lives? *Economic Inquiry*, 47(2), 317-336.

265 Nadler, J. (2020). H.R.3884 - Marijuana Opportunity Reinvestment and Expungement Act of
266 2019. Retrieved from <https://www.congress.gov/bill/116th-congress/house-bill/3884>

267 Naji, L., Rosic, T., Dennis, B., Bhatt, M., Sanger, N., Hudson, J., . . . Samaan, Z. (2018, June
268 11). The association between cannabis use and suicidal behavior in patients with
269 psychiatric disorders: an analysis of sex differences. *Biology of Sex Differences*, 9(22).

270 National Academies of Sciences, Engineering, and Medicine. (2017). *The Health Effects of
271 Cannabis and Cannabinoids: The Current State of Evidence and Recommendations for
272 Research*. Washington: National Academies Press.

273 National Institute on Alcohol Abuse and Alcoholism. (2020, April). Surveillance Report #115.
274 Washington, District of Columbia, United States.

275 Pederson, W. (2008, November). Does cannabis use lead to depression and suicidal behaviours?
276 A population-based longitudinal study. *Acta Psychiatrica Scandinavica*, 118(5), 395-403.

277 Robertson, L. S. (2014, December). Science and public health principles used to reduce road
278 deaths. *American Journal of Public Health*, 104(12), 2256-2258.

279 SAMHSA. (2015). *2013-2014 NSDUH State Estimates Of Substance Use And Mental Disorders*.
280 Rockville, MD.

281 SAMHSA. (2018). National Survey on Drug Use and Health (NSDUH-2017). Rockville,
282 Maryland, United States.

283 SAMHSA. (2019). *Key Substance Use and Mental Health Indicators in the United States:
284 Results from the 2018 National Survey on Drug Use and Health*. Rockville, MD:
285 Substance Abuse and Mental Health Services.

286 Tax Policy Center. (2020). State Alcohol Excise Taxes. Washington, District of Columbia,
287 United States.

288 Thornton, J. A. (2011, October 14). Does More Medical Care Improve Population Health? New
289 Evidence for an Old Controversy. *Applied Economics*, 34(1), 3325-3336.

290 Warshauer, M., & Monk, M. (1978, April). Problems in suicide statistics for whites and blacks.
291 *American Journal of Public Health*, 68(4), 383-388.

292

293

294

295

296

297

298

299

300

301

302

303

304 **Tables and Figures**

305


306 **Table 1 — Effective Dates of Marijuana Legalization**

State	FIPS	Recreational Effective	Medical Effective	Decriminalized Effective
Alabama	1			
Alaska	2	2/24/15	1999	1975-1990, 8/29/2003-6/14/2006, 2/24/2015
Arizona	4		3/14/11	
Arkansas	5		4/8/17	
California	6	11/9/16		1976
Colorado	8	1/3/13	6/1/01	1975
Connecticut	9		3/5/12	6/6/11
Delaware	10		5/13/11	12/18/15
District of Columbia	11	2/26/15	3/29/13	7/17/14
Florida	12		1/3/17	
Georgia	13			
Hawaii	15	1/11/20	8/8/17	1/11/20
Idaho	16			
Illinois	17	1/1/20	1/1/14	
Indiana	18			
Iowa	19			
Kansas	20			
Kentucky	21			
Louisiana	22		8/6/19	
Maine	23	12/17/16	5/10/10	5/1/76
Maryland	24		1/1/17	10/1/14
Massachusetts	25	12/15/16	1/1/13	1/2/09
Michigan	26	12/6/19	10/25/18	
Minnesota	27		7/1/15	1976
Mississippi	28			
Missouri	29		12/6/18	1/1/17
Montana	30		11/2/04	
Nebraska	31			1/1/79
Nevada	32	1/1/17	10/1/01	10/1/01
New Hampshire	33		7/23/13	9/16/17
New Jersey	34		1/1/11	
New Mexico	35		7/1/07	7/1/19
New York	36		7/7/14	7/29/77
North Carolina	37			7/1/77
North Dakota	38		1/28/17	5/2/19

Ohio	39		9/6/16	8/22/75
Oklahoma	40		8/30/18	
Oregon	41	7/1/15	12/3/98	10/1/73
Pennsylvania	42		5/17/16	
Rhode Island	44		1/3/06	4/1/13
South Carolina	45			
South Dakota	46			
Tennessee	47			
Texas	48			
Utah	49			
Vermont	50	7/1/18	5/19/04	7/1/13
Virginia	51			7/1/20
Washington	53	12/2/12	1998	12/2/12
West Virginia	54			
Wisconsin	55			
Wyoming	56			

307

308 Figure 1



309

Year of Law Change	Centred Years	Legal Slope	Illegal Slope
-3 to -1	2011-2013	0.7084	0.3744
0 to 3	2014-2017	0.3397	0.4116

310

311

312 Table 2 — Regression Results

Dependent Variable	Policies	Coefficients			Coefficients			Coefficients					
		Controls, TE, FE, ST	Conf.	[95% Interval]	Controls, TE, FE	Conf.	[95% Interval]	TE, FE	Conf.	[95% Interval]			
Total Suicide Rate ^a	Recreational Marijuana	-0.0299	*	-0.0567	-0.0030	-0.0444	-0.0918	0.0030	-0.0340	-0.0852	0.0171		
	Medical Marijuana	-0.0176		-0.0696	0.0344	0.0134	-0.0182	0.0450	0.0272	-0.0045	0.0590		
Male Suicide Rate ^a	Recreational Marijuana	0.0749		-0.1508	0.3006	0.0783	-0.0963	0.2529	0.0838	-0.0917	0.2593		
	Medical Marijuana	-0.0587		-0.1683	0.0510	-0.0073	-0.0761	0.0614	-0.0070	-0.0729	0.0590		
Female Suicide Rate ^a	Recreational Marijuana	-0.0373		-0.0938	0.0192	-0.0561	-0.1330	0.0207	-0.0432	-0.1275	0.0410		
	Medical Marijuana	0.0233		-0.0500	0.0967	0.0546	*	0.0091	0.1001	0.0766	**	0.0278	0.1253
Suicide Rate Male 15 to 19 ^b	Recreational Marijuana	-0.0843		-0.2344	0.0657	0.0319	-0.1021	0.1660	0.0387	-0.0756	0.1530		
	Medical Marijuana	0.0058		-0.0730	0.0846	-0.0263	-0.1093	0.0566	-0.0242	-0.1050	0.0566		
Suicide Rate Male 20 to 29 ^a	Recreational Marijuana	-0.0131		-0.0645	0.0382	-0.0018	-0.0589	0.0553	0.0142	-0.0368	0.0652		
	Medical Marijuana	-0.0084		-0.0755	0.0587	-0.0136	-0.0805	0.0534	-0.0200	-0.0852	0.0451		
Suicide Rate Male 30 to 39 ^a	Recreational Marijuana	-0.0026		-0.0422	0.0370	-0.0184	-0.0822	0.0454	-0.0214	-0.0834	0.0406		
	Medical Marijuana	-0.0213		-0.0739	0.0313	-0.0263	-0.0674	0.0148	-0.0252	-0.0627	0.0124		
Suicide Rate Male 40 to 49 ^a	Recreational Marijuana	-0.0543	**	-0.0942	-0.0145	-0.0873	**	-0.1400	-0.0346	-0.0824	**	-0.1402	-0.0245
	Medical Marijuana	-0.0339		-0.1186	0.0507	0.0110		-0.0406	0.0626	0.0307		-0.0231	0.0845
Suicide Rate Male 50 to 59 ^a	Recreational Marijuana	-0.0259		-0.0629	0.0110	-0.0973	**	-0.1535	-0.0412	-0.0827	*	-0.1567	-0.0086
	Medical Marijuana	-0.0357		-0.0840	0.0126	0.0195		-0.0213	0.0603	0.0439		-0.0059	0.0938
Suicide Rate Male 60 and Up ^a	Recreational Marijuana	0.0023		-0.0555	0.0602	-0.0545		-0.1126	0.0036	-0.0469		-0.1187	0.0250
	Medical Marijuana	-0.0216		-0.0826	0.0394	0.0219		-0.0133	0.0570	0.0340		-0.0043	0.0723
Suicide Rate Female 15 to 19 ^c	Recreational Marijuana	-0.1013		-0.2976	0.0950	-0.1706		-0.3420	0.0007	-0.1431		-0.3040	0.0178
	Medical Marijuana	-0.1753		-0.3846	0.0340	-0.0761		-0.2200	0.0678	-0.0839		-0.2142	0.0465
Suicide Rate Female 20 to 29 ^d	Recreational Marijuana	0.0057		-0.0920	0.1034	-0.0470		-0.1198	0.0257	-0.0281		-0.1033	0.0471
	Medical Marijuana	-0.0484		-0.1596	0.0627	-0.0167		-0.0909	0.0576	0.0056		-0.0674	0.0785
Suicide Rate Female 30 to 39 ^e	Recreational Marijuana	0.0121		-0.0686	0.0928	-0.0354		-0.1182	0.0474	-0.0185		-0.0950	0.0579
	Medical Marijuana	0.0900		-0.0005	0.1806	0.1128	**	0.0450	0.1807	0.1276	**	0.0554	0.1998

It is made available under a CC-BY 4.0 International license.

It is made available under a CC-BY 4.0 International license.

Suicide Rate Female 40 to 49 ^f	Recreational Marijuana	-0.0867	-0.1762	0.0028	-0.1547	**	-0.2558	-0.0537	-0.1234	*	-0.2249	-0.0219
	Medical Marijuana	0.0001	-0.0854	0.0856	0.0620		-0.0209	0.1450	0.0919		-0.0021	0.1859
Suicide Rate Female 50 to 59 ^b	Recreational Marijuana	-0.0532	-0.1900	0.0836	-0.0548		-0.2322	0.1227	-0.0767		-0.2684	0.1151
	Medical Marijuana	0.0750	-0.0257	0.1757	0.0808		-0.0079	0.1695	0.1019	*	0.0098	0.1940
Suicide Rate Female 60 and Up ^e	Recreational Marijuana	-0.0009	-0.1391	0.1373	0.0031		-0.1335	0.1396	-0.0051		-0.1464	0.1363
	Medical Marijuana	0.0388	-0.0764	0.1540	0.0396		-0.0363	0.1156	0.0580		-0.0157	0.1317
Reports of Major Depression 12 to 17 ^h	Recreational Marijuana	-0.0401	-0.0864	0.0063	0.0171		-0.0241	0.0583	0.0278		-0.0048	0.0604
	Medical Marijuana	0.0107	-0.0443	0.0657	0.0073		-0.0263	0.0409	0.0141		-0.0193	0.0476
Reports of Major Depression 18 to 25 ^h	Recreational Marijuana	0.0614	-0.0103	0.1331	0.0908	**	0.0515	0.1300	0.0927	**	0.0650	0.1205
	Medical Marijuana	0.0004	-0.0461	0.0468	0.0125		-0.0229	0.0478	0.0192		-0.0126	0.0510
Reports of Major Depression 26 and Up ^h	Recreational Marijuana	-0.0508	-0.1227	0.0210	0.0115		-0.0329	0.0559	0.0138		-0.0236	0.0512
	Medical Marijuana	-0.0084	-0.0409	0.0241	-0.0078		-0.0437	0.0281	-0.0027		-0.0348	0.0294
Reports of Major Depression 18 and Up ^h	Recreational Marijuana	-0.0220	-0.0852	0.0411	0.0315		-0.0071	0.0701	0.0357		0.0061	0.0654
	Medical Marijuana	-0.0066	-0.0380	0.0248	-0.0031		-0.0341	0.0278	0.0026		-0.0247	0.0299
Serious Mental Illness 18 to 25 ⁱ	Recreational Marijuana	0.0717	-0.0438	0.1872	0.0837	**	0.0375	0.1299	0.0603	**	0.0223	0.0983
	Medical Marijuana	-0.0186	-0.0802	0.0431	-0.0042		-0.0416	0.0333	-0.0035		-0.0445	0.0376
Serious Mental Illness 26 and Up ⁱ	Recreational Marijuana	-0.0658	-0.1630	0.0314	-0.0060		-0.0639	0.0518	-0.0023		-0.0486	0.0440
	Medical Marijuana	0.0454	-0.0379	0.1286	0.0024		-0.0370	0.0418	-0.0063		-0.0483	0.0357
Serious Mental Illness 18 and Up ⁱ	Recreational Marijuana	-0.0375	-0.1300	0.0550	0.0096		-0.0440	0.0632	0.0128		-0.0304	0.0561
	Medical Marijuana	0.0335	-0.0419	0.1090	0.0029		-0.0301	0.0358	-0.0032		-0.0388	0.0324
Serious Thoughts of Suicide 18 to 25 ⁱ	Recreational Marijuana	-0.0124	-0.0554	0.0306	0.0333		-0.0322	0.0988	0.0192		-0.0545	0.0929
	Medical Marijuana	0.0039	-0.0702	0.0780	-0.0240		-0.0648	0.0168	-0.0205		-0.0579	0.0168
Serious Thoughts of Suicide 26 and Up ⁱ	Recreational Marijuana	-0.0038	-0.0980	0.0905	0.0603		-0.0075	0.1280	0.0470		-0.0042	0.0983
	Medical Marijuana	0.0489	-0.0112	0.1091	-0.0281		-0.0790	0.0229	-0.0295		-0.0899	0.0310
Serious Thoughts of Suicide 18	Recreational Marijuana	-0.0034	-0.0789	0.0721	0.0464		-0.0148	0.1076	0.0342		-0.0157	0.0841

It is made available under a CC-BY 4.0 International license.

and Up ⁱ	Medical Marijuana	0.0330	-0.0157	0.0817	-0.0291	-0.0630	0.0048	-0.0285	-0.0719	0.0149
Any Mental Illness 18 to 25 ⁱ	Recreational Marijuana	-0.0052	-0.0452	0.0348	0.0260	-0.0019	0.0538	0.0341	**	0.0090
	Medical Marijuana	0.0012	-0.0429	0.0454	0.0101	-0.0168	0.0369	0.0092	-0.0170	0.0354
Any Mental Illness 26 and Up ⁱ	Recreational Marijuana	-0.0277	-0.0898	0.0345	0.0295	-0.0135	0.0725	0.0230	-0.0127	0.0588
	Medical Marijuana	-0.0146	-0.0598	0.0307	-0.0248	-0.0710	0.0213	-0.0269	-0.0738	0.0199
Any Mental Illness 18 and Up ⁱ	Recreational Marijuana	-0.0230	-0.0776	0.0315	0.0286	-0.0087	0.0659	0.0258	-0.0049	0.0566
	Medical Marijuana	-0.0119	-0.0519	0.0280	-0.0191	-0.0590	0.0209	-0.0204	-0.0605	0.0198

313 * $P < .05$

314 ** $P < .01$

315 ^a n = 1 020

316 ^b n = 1 012

317 ^c n = 951

318 ^d n = 1 005

319 ^e n = 1 008

320 ^f n = 1 014

321 ^g n = 1 009

322 ^h n = 663

323 ⁱ n = 510